

FACT SHEET NO. 14

# PHYSICAL ACTIVITY LEVELS OF THE GREAT SOUTHERN REGION

## The Premier's Physical Activity Taskforce

In November 2002, the Premier's Physical Activity Taskforce (PATF) undertook a state-wide survey to measure physical activity levels of Western Australian adults. This fact sheet presents major findings from this survey for the Great Southern region<sup>1</sup>.

## Great Southern Sample Demographics

The Great Southern region has an estimated population of 27,060 (in 2001). From the Great Southern region, 69 participants or 2.2% of the total sample (n = 3,200) completed the physical activity survey.

Table 1: Demographic Profile of Respondents from the Great Southern Region


| CHARACTERISTIC   | %    |
|------------------|------|
| <b>Gender</b>    |      |
| Male             | 53.6 |
| Female           | 46.4 |
| <b>Age Group</b> |      |
| 18 to 29 years   | 11.6 |
| 30 to 44 years   | 26.1 |
| 45 to 59 years   | 42.0 |
| 60 years or more | 20.3 |
| <b>Education</b> |      |
| Less than TEE    | 50.7 |
| TEE/diploma      | 36.2 |
| University       | 13.0 |

(Note n = 69)

## Levels of Sufficient Physical Activity

Figure 1 presents the proportion of adults from the Great Southern region who were:


- **Sufficiently active** ( $\geq 150$  minutes of moderate activity over 5 or more days or  $\geq 60$  minutes of vigorous activity per week);
- **Insufficiently active** (some physical activity but less than sufficient), and;
- **Inactive** (no physical activity).



The proportion of Great Southern adults who were sufficiently active was less (52.2%) than Western Australian adults overall (54.6%). The proportion of Great Southern adults who were inactive was higher (18.8%) than Western Australian adults overall (13.5%).

## Levels of Sufficient Physical Activity by Gender

Figure 2 presents the proportion of Great Southern (GS) and Western Australian (WA) adults who were inactive and sufficiently active by gender.




Great Southern males were more inactive (23.4%) and less sufficiently active (50%) compared to females from the Great Southern region and Western Australian adults overall.



### Participation in Sufficient Moderate and Vigorous Physical Activity by Gender

Figure 3 presents the proportion of Great Southern adults participating in sufficient moderate ( $\geq 150$  minutes of moderate-intensity physical activity over 5 or more days per week) and vigorous ( $\geq 60$  minutes of vigorous-intensity physical activity per week) physical activity.

Figure 3. Levels of Sufficient Moderate and Vigorous Physical Activity



A lower proportion of Great Southern males (23.5%) and a higher proportion of females (45.7%) participated in sufficient levels of moderate physical activity compared to Western Australian adult males (35.4%) and females (37.7%) overall. A lower proportion of Great Southern males (34.3%) and females (17.1%) participated in sufficient levels of vigorous physical activity compared to Western Australian adult males (37%) and females (25.8%) overall.

### Other Physical Activity Facts for the Great Southern Region (WA results in brackets)

- 40.6% (62%) walked for recreation or exercise, 20.3% (28%) walked for transport, and 4.3% (8%) cycled for recreation in the week before the survey.
- 38.1% (57%) used the streets or footpaths, and 6.3% (17%) used a public park or oval for physical activity in the week before the survey.
- 56.5% (49%) reported “not having enough time” as a barrier to being more active and 36.9% (34%) reported “to improve fitness” as a major reason for participating in physical activity.




### Overweight and Obesity

Self-reported height and weight data were used to determine the Body Mass Index (BMI) for each survey participant (weight in kilograms divided by height in metres squared). The following BMI categories were used to define weight<sup>2</sup>:

- $< 18.50$  (underweight)
- 18.50-24.99 (healthy weight)
- 25.00-29.99 (overweight)
- $\geq 30.00$  (obese)

The prevalence for each of these BMI categories in the Great Southern region is presented in Figure 4.

Figure 4. Levels of BMI



A lower proportion of Great Southern adults were in the healthy weight range (36.2%) and higher proportions were overweight (43.5%) and obese (18.8%) compared to the overall Western Australian adult population (healthy weight 49.8% overweight 34.7%, and obese 13%).

### Recommendations

Recommendations for increasing levels of sufficient physical activity in Western Australia, as well as key findings can be found in the recently published report, Physical Activity Levels of Western Australian Adults 2002<sup>3</sup>. This report can be obtained from the PATF website ([www.patf.dpc.wa.gov.au](http://www.patf.dpc.wa.gov.au)) or by contacting the PATF Secretariat on 9382 5980.

© Government of Western Australia 2003

There is no objection to this publication being copied in whole or in part, provided there is due acknowledgement of any material quoted from the publication.

**Suggested Citation:** McCormack, G., & Watt, S. (2003). *Physical Activity Levels of the Great Southern Region (Fact Sheet No.14)*. Perth, Western Australia: Western Australian Government.



Department of Health  
Department of Sport  
and Recreation



<sup>1</sup> As defined by the Department of Sport and Recreation, WA

<sup>2</sup> World Health Organisation (2001). Obesity: Preventing and Managing the Global Epidemic. WHO Technical Series: No. 894. Geneva: WHO; 2000.

<sup>3</sup> McCormack G, Milligan R, Giles-Corti B, & Clarkson J. (2003). Physical Activity Levels of Western Australian Adults 2002. Results from the Adult Physical Activity Survey and Pedometer Study. Perth, Western Australia: Western Australian Government.